

Quotient Star Bodies, Intersection Bodies, and Star Duality

Maria Moszyńska*

metadata, citation and similar papers at core.ac.uk

Submitted by Ying-ming Liu

Received March 25, 1998

INTRODUCTION

Various topological constructions have their metric counterparts; for example, metric product corresponds to topological product (see [6]) and metric inverse limit corresponds to topological inverse limit (see [9, 10]). The main subject of this paper is a metric counterpart of the topological notion of quotient space (see [2] or [3]). We restrict our consideration to a compact metric space A and a very simple equivalence relation, with only one nontrivial equivalence class $C \subset A$. Then, evidently, the natural quotient map $p: A \rightarrow A/C$ (defined by $p(x) = [x]$) satisfies the condition

$$p(C) \text{ is a singleton and } p|(A \setminus C) \text{ is a topological embedding.} \quad (0.1)$$

Topologists used to look at quotient space “up to a homeomorphism”; i.e., more precisely, they work with the category *Top* of topological spaces and continuous maps. For geometers, it is important to choose a “good” representative of the topological type of A/C ; this representative will be the image of A under a map p_C satisfying (0.1), with some nice geometric properties.

We shall keep in mind the following physical interpretation. Let A be a soft body with a stone C inside. When the stone is cut out, the hole shrinks to a point and the remaining part $A \setminus C$ changes its geometric shape. The shape of the resulting body A/C depends not only on geometric properties of A and C , but also on physical properties of A and forces acting on A .

* E-mail: mariamos@mimuw.edu.pl.

This interpretation justifies further restrictions. We assume that A is a subset of \mathbb{R}^n , $0 \in C$, and $A \setminus C$ is homeomorphic to $A \setminus \{0\}$. Then A/C is homeomorphic to a subset of \mathbb{R}^n and so we can look for a map p_C satisfying (0.1), with values in \mathbb{R}^n . Moreover, if $p_C: \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfying (0.1) is independent of A , then it is reasonable to consider properties of A invariant under p_C .

Following [3], we shall refer to p_C as a *quotient map* (an *identification* in terminology of [2]).

The following theorem follows directly from [2, Theorem 7.7, p. 17 and Theorem 4.3, p. 126].

THEOREM 0.1. *Let \mathcal{C} be a nonempty family of compact subsets of \mathbb{R}^n with $0 \in C$ and $\mathbb{R}^n \setminus C$ homeomorphic to $\mathbb{R}^n \setminus \{0\}$ for every $C \in \mathcal{C}$, and let p_C be a quotient map with $p_C(C) = 0$. For every map $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ preserving \mathcal{C} , with $f(0) = 0$, and for every $C \in \mathcal{C}$, let*

$$\hat{f}(x) := p_{f(C)} f p_C^{-1}(x) \quad \text{for every } x \in \mathbb{R}^n. \quad (0.2)$$

Then

(i) *the formula (0.2) defines a map $\hat{f}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $\hat{f}(0) = 0$ and the diagram*

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{f} & \mathbb{R}^n \\ p_C \downarrow & & \downarrow p_{f(C)} \\ \mathbb{R}^n & \xrightarrow{\hat{f}} & \mathbb{R}^n \end{array}$$

commutes, and

(ii) *if f is a homeomorphism, then so is \hat{f} .*

We shall work with the class of star bodies in \mathbb{R}^n with nonnegative and continuous radial maps. Section 1 concerns star bodies and star maps. In Section 2 we define radial quotient maps and study corresponding quotient bodies. In particular, we prove that the operation $(A, C) \mapsto A/C$ preserves the class of intersection bodies of star bodies (Corollary 2.8).

Section 3 deals with the category \mathbf{St}^n of star bodies and star maps, and its subcategory \mathbf{St}_+^n with star bodies whose radial functions are positive. We define and study a duality on this subcategory: a functor from \mathbf{St}_+^n to itself which is an involution (compare [11]).

We use the following notation: For affine independent points x_0, \dots, x_k in \mathbb{R}^n , the simplex with vertices x_0, \dots, x_k is denoted by $\Delta(x_0, \dots, x_k)$; in particular, if a, b are distinct points, then $\Delta(a, b)$ is the segment with endpoints a, b .

As usual, B^n and S^{n-1} are the unit ball and the unit sphere, and κ_n is the volume of B^n . We denote by σ the spherical Lebesgue measure on S^{n-1} . For any $u \in S^{n-1}$, the (linear) hyperplane orthogonal to u is denoted by u^\perp . For any convex body $A \subset \mathbb{R}^n$, its support function is $h(A, \cdot)$, supporting hyperplane is $H(A, \cdot)$, and $b(A)$ is the mean width of A . The class of convex bodies in \mathbb{R}^n is \mathcal{K}_0^n .

For basic notions of the category theory, see [8].

1. STAR BODIES AND STAR MAPS

Since in the literature there are various notions of a star body (see [4, p. 18; [12], p. 416), we start from definitions: Let A be a nonempty compact subset of \mathbb{R}^n and $a \in A$. Then, A is a *body* if and only if $A = \text{cl int } A$, and A is *star shaped at a* if and only if $\Delta(a, x) \subset A$ for every $x \in A$.

For convenience, we shall assume that $a = 0$. The *radial function* $\varrho_A: \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{R}$ of a set A which is star shaped at 0 is defined by the formula

$$\varrho_A(x) := \sup\{\lambda \geq 0 \mid \lambda x \in A\}.$$

(Sometimes, we write $\varrho(A, x)$ instead of $\varrho_A(x)$.) Obviously, ϱ_A is homogeneous of degree -1 .

A set $A \subset \mathbb{R}^n$ will be called a *star body* whenever A is a body which is star shaped at 0 and whose radial function restricted to S^{n-1} is continuous.

Let us notice that this notion of star body is less general than that of Gardner [4], since we assume that $0 \in A$ and thus $\varrho_A \geq 0$. Moreover, Gardner assumes only continuity of the restriction of radial function to its support. On the other hand, our notion is more general than that of Schneider [12], since we do not require $\varrho_A > 0$; that is, we allow 0 to belong to $\text{bd } A$.

Let \mathcal{S}_0^n be the class of all the star bodies in \mathbb{R}^n and let

$$\mathcal{S}_1^n = \mathcal{S}_0^n \cup \{\{0\}\}.$$

We are looking for a possibly large group of transformation which preserve \mathcal{S}_0^n and \mathcal{S}_1^n .

DEFINITION 1.1. A map $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a *star map* if and only if f is a positively homogeneous homeomorphism.

Of course, the set of all the star maps of \mathbb{R}^n is a group of transformations. We shall denote it by $\text{GS}(n)$ (general star maps).

The following is evident.

PROPOSITION 1.2. *If $f \in \text{GS}(n)$ and $A \in \mathcal{S}_0^n$, then for every $x \in \mathbb{R}^n$,*

$$\varrho_{f(A)}(f(x)) = \varrho_A(x).$$

The next statement is a direct consequence of Proposition 1.2.

PROPOSITION 1.3. *The classes \mathcal{S}_0^n and \mathcal{S}_1^n are invariant under star maps.*

Let us prove the following.

PROPOSITION 1.4. *Every two star bodies A_1, A_2 with 0 in the interior are star equivalent; i.e., there exists $g \in \text{GS}(n)$ such that $g(A_1) = A_2$.*

Proof. Let $A_1, A_2 \in \mathcal{S}_0^n$. We define $g: \mathbb{R}^n \rightarrow \mathbb{R}^n$ by the formula

$$g(x) = \begin{cases} \frac{\varrho_{A_2}(x)}{\varrho_{A_1}(x)} \cdot x, & \text{if } x \neq 0, \\ 0, & \text{if } x = 0. \end{cases} \quad (1.1)$$

It is clear that $g \in \text{GS}(n)$.

Let us show that $g(A_1) = A_2$. Indeed, g is a homeomorphism and maps $\text{bd } A_1$ onto $\text{bd } A_2$:

$$\begin{aligned} x \in \text{bd } A_1 &\Leftrightarrow \varrho_{A_1}(x) = 1 \Leftrightarrow g(x) = \varrho_{A_2}(x) \cdot x \\ &\Leftrightarrow \varrho_{A_2}(g(x)) = 1 \Leftrightarrow g(x) \in \text{bd } A_2. \end{aligned}$$

■

Evidently, $\text{GL}(n)$ is a subgroup of $\text{GS}(n)$. As a consequence of Proposition 1.4, we obtain the following corollary:

COROLLARY 1.5. *$\text{GL}(n)$ is a proper subgroup of $\text{GS}(n)$.*

Let us notice that the group $\text{GS}(n)$ is generated by its two subgroups,

$$\text{SS}(n) := \{f \in \text{GS}(n) \mid f(S^{n-1}) = S^{n-1}\} \quad (1.2)$$

and

$$\text{RS}(n) := \{f \in \text{GS}(n) \mid \exists \lambda: \mathbb{R}^n \rightarrow \mathbb{R}_+, f(x) = \lambda(x) \cdot x \text{ for every } x\}. \quad (1.3)$$

The first subgroup consists of *spherical star maps*, the star extensions of homeomorphisms of S^{n-1} onto itself; the second subgroup consists of *radial star maps*, the star extensions of central projections of S^{n-1} .

PROPOSITION 1.6. *For every $f \in \text{GS}(n)$ there exists $g \in \text{RS}(n)$ and $h \in \text{SS}(n)$ such that $f = gh$. This decomposition is unique.*

Proof. Let $f \in \text{GS}(n)$ and let $A = f(B^n)$. Then A is a star body with $0 \in \text{int } A$, and thus, by Proposition 1.4, there exists $g \in \text{GS}(n)$ such that $g(S^{n-1}) = \text{bd } A$. The map g is defined by (1.1) for $A_1 = B^n$ and $A_2 = A$, whence $g \in \text{RS}(n)$.

Let

$$h := g^{-1}f.$$

Then h is a star map preserving S^{n-1} , and $f = gh$.

To prove the uniqueness of the decomposition of f , suppose that

$$f = g_1 h_1 = g_2 h_2$$

for some radial star maps g_1, g_2 and spherical star maps h_1, h_2 . Then $g_2^{-1}g_1 = h_2 h_1^{-1}$, where the left side belongs to $\text{RS}(n)$ and the right side belongs to $\text{SS}(n)$. Since these two subgroups have only the identity map in common, it follows that $g_1 = g_2$ and $h_1 = h_2$. ■

Every map $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ preserving \mathcal{S}_1^n induces the map $f_*: \mathcal{S}_1^n \rightarrow \mathcal{S}_1^n$ defined by

$$f_*: A \mapsto f(A).$$

According to [4], the radial metric δ in \mathcal{S}_1^n is defined by the formula

$$\delta(A_1, A_2) = \sup_{u \in S^{n-1}} |\varrho_{A_1}(u) - \varrho_{A_2}(u)|.$$

THEOREM 1.7. *If $f \in \text{GS}(n)$, then f_* is a Lipschitz map with respect to the radial metric δ . If, moreover, $f \in \text{SS}(n)$, then f_* is an isometry.*

Proof. Let $A_1, A_2 \in \mathcal{S}_1^n$. Then, by Proposition 1.2,

$$\begin{aligned} & \delta(f_*(A_1), f_*(A_2)) \\ &= \sup_{u \in S^{n-1}} |\varrho_{f(A_1)}(u) - \varrho_{f(A_2)}(u)| \\ &= \sup_{u \in S^{n-1}} |\varrho_{A_1}(f^{-1}(u)) - \varrho_{A_2}(f^{-1}(u))| \\ &= \sup_u \frac{1}{\|f^{-1}(u)\|} \left| \varrho_{A_1}\left(\frac{f^{-1}(u)}{\|f^{-1}(u)\|}\right) - \varrho_{A_2}\left(\frac{f^{-1}(u)}{\|f^{-1}(u)\|}\right) \right| \\ &\leq \sup \frac{1}{\|f^{-1}(u)\|} \cdot \delta(A_1, A_2), \end{aligned}$$

where $\sup(1/\|f^{-1}(u)\|) < \infty$, because f^{-1} is a bijection, $f(0) = 0$, and S^{n-1} is compact.

The second assertion is obvious. ■

2. RADIAL QUOTIENT STAR BODIES

We shall consider the following family of maps of \mathbb{R}^n onto itself.

DEFINITION 2.1. Let $C \in \mathcal{S}_1^n$. We define $p_C: \mathbb{R}^n \rightarrow \mathbb{R}^n$ by the formula

$$p_C(x) := \begin{cases} 0, & \text{if } x \in C, \\ (1 - \varrho_C(x)) \cdot x, & \text{if } x \in \mathbb{R}^n \setminus \text{int } C. \end{cases}$$

PROPOSITION 2.2. For every $C \in \mathcal{S}_1^n$:

- (i) p_C is a quotient map;
- (ii) if A is a star body containing C and $A \neq C$, then $p_C(A) \in \mathcal{S}_0^n$.

In view of Proposition 2.2, we shall refer to p_C as a *radial quotient map* and to $p_C(A)$ as a *radial quotient star body* of A . We shall use the symbol A/C also for $p_C(A)$.

Let us note the following.

PROPOSITION 2.3. If $A, C \in \mathcal{S}_1^n$ and $C \subset A$, then

$$\varrho(A/C, x) = \varrho(A, x) - \varrho(C, x)$$

for every $x \in \mathbb{R}^n \setminus \{0\}$.

Proof. It suffices to prove the assertion for $\varrho|_{S^{n-1}}$. Let $u \in S^{n-1}$ and $x_0 \in \text{bd } A \cap \text{pos } u$. Then

$$\begin{aligned} \varrho(p_C(A), u) &= \|p_C(x_0)\| = (1 - \varrho_C(x_0)) \cdot \|x_0\| \\ &= \|x_0\| - \varrho_C\left(\frac{x_0}{\|x_0\|}\right) = \varrho_A(u) - \varrho_C(u). \end{aligned}$$

■

The following statement is a direct consequence of Proposition 2.3.

COROLLARY 2.4. The function $\Phi: \{(A, C) \in \mathcal{S}_1^n \times \mathcal{S}_1^n \mid C \subset A\} \rightarrow \mathcal{S}_1^n$ defined by

$$\Phi(A, C) = A/C \tag{2.1}$$

is continuous with respect to the radial metric δ .

For a given $C \in \mathcal{S}_1^n$, let

$$\mathcal{S}_C^n := \{A \in \mathcal{S}_1^n \mid A \supset C\}. \tag{2.2}$$

THEOREM 2.5. For every $C \in \mathcal{S}_1^n$, there exists a homotopy

$$(\psi_t: \mathbb{R}^n \rightarrow \mathbb{R}^n)_{t \in [0, 1]}$$

such that $\psi_0 = \text{id}_{\mathbb{R}^n}$, $\psi_1 = p_C$, and for every $t \in [0, 1]$, the induced map $(\psi_t)_*: \mathcal{S}_C^n \rightarrow \mathcal{S}_1^n$ is an isometry with respect to δ .

Proof. We define $\psi_t: \mathbb{R}^n \rightarrow \mathbb{R}^n$ by the formula

$$\psi_t(x) = \begin{cases} (1-t) \cdot x, & \text{if } x \in C, \\ (1-t\varrho_C(x)) \cdot x, & \text{if } x \notin C. \end{cases}$$

Evidently, $\psi: \mathbb{R}^n \times [0, 1] \ni (x, t) \mapsto \psi_t(x) \in \mathbb{R}^n$ is continuous,

$$\psi(x, 0) = x, \quad \text{and} \quad \psi(x, 1) = p_C(x).$$

Let $t \in [0, 1]$ and $A_1, A_2 \in \mathcal{S}_C^n$. We shall show that

$$\delta(A_1, A_2) = \delta(\psi_t(A_1), \psi_t(A_2)). \quad (2.3)$$

If $u \in S^{n-1}$, $i \in \{1, 2\}$, and $a_i \in \text{bd } A_i \cap \text{pos } u$, then $\varrho(A_i, u) = \|a_i\|$ and

$$\varrho(\psi_t(A_i), u) = \|\psi_t(a_i)\| = \begin{cases} (1-t)\|a_i\|, & \text{if } a_i \in C, \\ (1-t\varrho_C(a_i))\|a_i\|, & \text{if } a_i \notin C. \end{cases}$$

Since $(1-t\varrho_C(a_i))\|a_i\| = \|a_i\| - t\varrho_C(u)$ and $a_1, a_2 \in C$ implies $a_1 = a_2$, by easy calculation it follows that

$$|\varrho(\psi_t(A_1), u) - \varrho(\psi_t(A_2), u)| = \begin{cases} 0, & \text{if } a_1, a_2 \in C, \\ \left| \|a_1\| - \|a_2\| \right|, & \text{otherwise.} \end{cases}$$

Thus,

$$\delta(\psi_t(A_1), \psi_t(A_2)) = \sup_{u \in S^{n-1}} |\varrho(A_1, u) - \varrho(A_2, u)| = \delta(A_1, A_2),$$

which proves (2.3). ■

Let us observe that for every star map f the induced map f_* is a homomorphism with respect to the operation Φ defined by (2.1):

PROPOSITION 2.6. *For every $f \in \text{GS}(n)$ and $A, C \in \mathcal{S}_1^n$ with $C \subset A$,*

$$f(A/C) = f(A)/f(C).$$

Proof. Since f is positively homogeneous and $\varrho_{f(C)}(f(x)) = \varrho_C(x)$, by Definition 2.1 it follows that

$$fp_C(x) = p_{f(C)}(f(x)) \quad \text{for every } x \in \mathbb{R}^n.$$

■

The operation Φ is also invariant under the function $I_1: \mathcal{S}_1^n \rightarrow \mathcal{S}_1^n$, the

first of I_1, \dots, I_{n-1} , which are defined as follows: For every $u \in S^{n-1}$,

$$\varrho(I_k A, u) := \tilde{V}_{k, n-1}(A \cap u^\perp),$$

where

$$\tilde{V}_{k, n-1}(A \cap u^\perp) = \frac{1}{n-1} \int_{S^{n-1} \cap u^\perp} \varrho(A, v)^k d\sigma(v)$$

for any $k \in \{1, \dots, n-1\}$ (see [4] or [7]). If $A \in \mathcal{S}_0^n$, then $I_k A$ is called the *intersection body of order k* of the star body A .

THEOREM 2.7. *For every $A, C \in \mathcal{S}_1^n$ with $C \subset A$,*

$$I_1(A/C) = I_1 A / I_1 C.$$

Proof. By Proposition 2.3, for every affine subspace E in \mathbb{R}^n ,

$$(A/C) \cap E = (A \cap E) / (C \cap E). \quad (2.4)$$

Let $u \in S^{n-1}$. Using Proposition 2.3 combined with (2.4) for $E = u^\perp$, we obtain

$$\varrho(I_1(A/C), u) = \varrho(I_1 A / I_1 C, u).$$

■

As was noticed in [4, Note 8.3, p. 305], for any $k \in \{1, \dots, n-1\}$, a set A is the intersection body of order k of a star body if and only if A is the intersection body of a star body. Thus Theorem 2.7 yields the following corollary:

COROLLARY 2.8. *Let $A, C \in \mathcal{S}_1^n$ and $C \subset A$. If A and C are the intersection bodies of some star bodies, then so is A/C .*

Corollary 2.8 can be useful in geometric tomography, where the intersection bodies play an essential role. See, for instance, Theorem 8.2.8 of [4], which gives an answer to the question whether

$$V_{n-1}(A_1 \cap u^\perp) \leq V_{n-1}(A_2 \cap u^\perp) \quad \text{for every } u \in S^{n-1}$$

implies $V_n(A_1) \leq V_n(A_2)$. According to that theorem, the answer is positive when A_1 is the intersection body of a star body.

3. CATEGORICAL APPROACH. DUALITY OF STAR BODIES

Let \mathbf{St}^n be the category whose objects are star bodies in \mathbb{R}^n :

$$\text{Ob } \mathbf{St}^n = \mathcal{S}_0^n,$$

and for every $A_1, A_2 \in \mathcal{S}_0^n$ the set of morphisms, $\mathbf{St}^n(A_1, A_2)$, consists of star maps sending A_1 to A_2 . Furthermore, let \mathbf{St}_2^n be the category with the class of objects

$$\text{Ob } \mathbf{St}_2^n = \{(A, C) \in \mathcal{S}_0^n \times \mathcal{S}_0^n \mid C \subset A \neq C\}$$

and with morphisms being suitable star maps. The function $\Phi: \text{Ob } \mathbf{St}_2^n \rightarrow \mathcal{S}_0^n$ defined by (2.1) can be extended to a functor:

PROPOSITION 3.1. *The formulae*

$$\Phi(A, C) := A/C \quad \text{for } A, C \in \mathcal{S}_0^n$$

and

$$\Phi(f) := \hat{f} \quad \text{for } f \in \mathbf{St}_2^n((A_1, C_1), (A_2, C_2))$$

[see (0.2)] define a covariant functor $\Phi: \mathbf{St}_2^n \rightarrow \mathbf{St}^n$, such that any morphism and its image under Φ are restrictions of the same star map.

Proof. In view of Theorem 0.1, it suffices to show that $\hat{f} = f$ for every $f \in \text{GS}(n)$. It is easy to check that, for every $x \in \mathbb{R}^n \setminus \{0\}$,

$$p_C^{-1}(x) = (1 + \varrho_C(x)) \cdot x$$

and

$$p_C(\beta x) = (\beta - \varrho_C(x)) \cdot x$$

for every $\beta \in \mathbb{R}$. Thus,

$$\begin{aligned} \hat{f}(x) &= p_{f(C)} p_C^{-1}(x) = p_{f(C)} f((1 + \varrho_C(x)) \cdot x) \\ &= p_{f(C)} ((1 + \varrho_C(x)) f(x)) = (1 + \varrho_C(x) - \varrho_{f(C)}(f(x))) f(x) \\ &= f(x). \end{aligned}$$

■

Given a category \mathbf{C} of star bodies, it is natural to look for a *duality* in this category, that is, a functor from \mathbf{C} to itself being an involution (compare [11]). In what follows we shall restrict our consideration to star

bodies with positive radial function (i.e., with the origin in the interior). Let \mathbf{St}_+^n be the corresponding full subcategory of \mathbf{St}^n .

Let i be the inversion of the one-point compactification $\bar{\mathbb{R}}^n$ of \mathbb{R}^n , with respect to S^{n-1} :

$$i(x) := \frac{x}{\|x\|^2} \quad \text{for } x \in \mathbb{R}^n \setminus \{0\}. \quad (3.1)$$

DEFINITION 3.2. For every object A of \mathbf{St}_+^n ,

$$A^\circ = \text{cl}(\mathbb{R}^n \setminus i(A));$$

for every $f \in \text{GS}(n)$,

$$f^\circ := ifi|_{\mathbb{R}^n}.$$

Let us note the following.

PROPOSITION 3.3. Let $A \in \mathcal{S}_0^n$ and $0 \in \text{int } A$. Then $A^\circ \in \mathcal{S}_0^n$ and for every $u \in S^{n-1}$,

$$\varrho(A^\circ, u) = \frac{1}{\varrho(A, u)}.$$

Proof. By Definition 3.2, A° is a star body and

$$\varrho(A^\circ, u) = \|a\|,$$

where $a \in \text{bd}(\mathbb{R}^n \setminus i(A)) \cap \text{pos } u$. Since $\text{bd}(\mathbb{R}^n \setminus A) = i(\text{bd } A)$, it follows that $i(a) \in \text{bd } A \cap \text{pos } u$, and thus

$$\|i(a)\| = \varrho(A, u).$$

By (3.1), this completes the proof. ■

THEOREM 3.4. The pair of functions $A \mapsto A^\circ$, $f \mapsto f^\circ$ is a covariant functor from \mathbf{St}_+^n to itself such that

$$A^{\circ\circ} = A \quad \text{and} \quad f^{\circ\circ} = f \quad (3.2)$$

for every A and f .

Moreover,

$$A \subset B \quad \Rightarrow \quad B^\circ \subset A^\circ. \quad (3.3)$$

Proof. By Proposition 3.3, if A is a star body with 0 in the interior, then so is A° . Let $f \in \text{GS}(n)$ and $f(A) = B$. Then

$$f^\circ \in \text{GS}(n), \quad (3.4)$$

because f° is a homeomorphism of \mathbb{R}^n and for every positive α ,

$$f^\circ(\alpha x) = ifi(\alpha x) = f\left(\frac{1}{\alpha}i(x)\right) = \alpha ifi(x) = \alpha f^\circ(x).$$

Moreover,

$$f^\circ(A^\circ) = B^\circ, \quad (3.5)$$

because

$$f^\circ(A^\circ) = ifi(\text{cl}(\mathbb{R}^n \setminus i(A))) = \text{cl}(\mathbb{R}^n \setminus if(A)) = (f(A))^\circ.$$

By (3.4) and (3.5), we have a covariant functor.

Furthermore,

$$\begin{aligned} A^\circ &= \text{cl}(\mathbb{R}^n \setminus i(A^\circ)) = \text{cl}(\mathbb{R}^n \setminus i(\text{cl}(\mathbb{R}^n \setminus i(A)))) \\ &= \mathbb{R}^n \setminus i(\text{int } \text{cl}(\mathbb{R}^n \setminus i(A))) = A, \end{aligned}$$

since f and i are homeomorphisms, $0 \in \text{int } A$, and i is an involution. Evidently, $f^{\circ\circ} = f$. This proves (3.2). By similar arguments, (3.3) holds. ■

By Theorem 3.4, the pair of functions $A \mapsto A^\circ$, $f \mapsto f^\circ$ is a duality. We shall refer to this functor as the *star duality*.

For a convex body A with the origin in the interior, there is a simple connection between polar dual A^* and star dual A° .

THEOREM 3.5. *If $A \in \mathcal{K}_0^n$ and $0 \in \text{int } A$, then*

$$\tilde{V}_1(A^\circ) = \frac{\kappa_n}{2} \cdot b(A^*).$$

Proof. By Proposition 3.4 and Remark 1.7.7 in [12],

$$\varrho(A^\circ, u) = h(A^*, u) \quad \text{for every } u \in S^{n-1}. \quad (3.6)$$

Hence, by the formula (1.7.2) in [12],

$$\begin{aligned} \tilde{V}_1(A^\circ) &= \frac{1}{n} \int_{S^{n-1}} \varrho(A^\circ, u) d\sigma(u) = \frac{1}{n} \int_{S^{n-1}} h(A^*, u) d\sigma(u) \\ &= \frac{\kappa_n}{2} \cdot b(A^*). \end{aligned}$$

■

Generally, star dual of a convex body is different from its polar dual; see Fig. 1.




FIG. 1. A , big square; A^* , small square; A° , flower.

THEOREM 3.6. *Let $n \geq 2$. For every $A \in \mathcal{H}_0^n$ with $0 \in \text{int } A$, the following are equivalent:*

- (i) $A^\circ = A^*$;
- (ii) A is a centered ball.

Proof. (ii) \Rightarrow (i) is obvious.

Assume (i). Then, by (3.6),

$$\varrho(A^*, u) = h(A^*, u) \quad \text{for every } u \in S^{n-1}$$

and thus

$$\text{pos } u \cap \text{bd } A^* \subset H(A^*, u).$$

As a consequence, $\text{bd } A^*$ is of the class C^1 .

Since $E \cap A^\circ = (E \cap A)^\circ$ for every two-dimensional linear subspace E , without any loss of generality we may assume that $n = 2$.

Let $r: \mathbb{R} \rightarrow \text{bd } A^*$ be the parametrization of $\text{bd } A^*$ in polar coordinates,

$$r(t) = \bar{\varrho}(t) \cdot u(t),$$

where $\bar{\varrho}(t) = \varrho(A^*, u(t))$ and $u(t) = (\cos t, \sin t)$. Then $\langle r(t), r'(t) \rangle = 0$ for every t ; i.e., $\bar{\varrho} \cdot \bar{\varrho}' = 0$. Hence $\bar{\varrho}' = 0$ and thus ϱ_{A^*} is constant, and so is ϱ_A . This completes the proof. ■

The following statement is an analogue of the conjecture (7.4.33) in [12], concerning the polar duality of convex bodies.

THEOREM 3.7. *Let A be a star body in \mathbb{R}^n with $0 \in \text{int } A$. For any $p, q > 0$ with $1/p + 1/q = 1$,*

$$\tilde{V}_p(A^\circ) \cdot \tilde{V}_q(A)^{p-1} \geq \kappa_n^p. \quad (3.7)$$

In particular,

$$\tilde{V}_2(A^\circ) \cdot \tilde{V}_2(A) \geq \kappa_n^2. \quad (3.8)$$

The equality in (3.7) or (3.8) holds if and only if A is a centered ball.

Proof. Since $\varrho_{A^\circ} \cdot \varrho_A = 1$ (see Proposition 3.3), by the Hölder inequality,

$$\begin{aligned} n \kappa_n &= \sigma(S^{n-1}) = \int_{S^{n-1}} \varrho_{A^\circ} \varrho_A \, d\sigma \\ &\leq \left(\int_{S^{n-1}} (\varrho_{A^\circ})^p \, d\sigma \right)^{1/p} \cdot \left(\int_{S^{n-1}} (\varrho_A)^q \, d\sigma \right)^{1/q} \\ &= (n \tilde{V}_p(A^\circ))^{1/p} \cdot (n \tilde{V}_q(A))^{1/q} = n \tilde{V}_p(A^\circ)^{1/p} \cdot \tilde{V}_q(A)^{(p-1)/p}. \end{aligned}$$

This implies (3.7) (and 3.8).

The equality holds if and only if there exists $\alpha > 0$ such that $\varrho_{A^\circ} = \alpha \varrho_A$. This happens if and only if A is a centered ball. ■

Finally, let us return to the notion of intersection body of order k . It has been defined only for $k > 0$ (see [4, Note 8.3]), but its definition can be automatically extended over $k < 0$. The following statement is evident:

3.8. For every star body A with $0 \in \text{int } A$,

$$I_{-k} A = I_k(A^\circ).$$

There is a relationship between star duality and the operation I_k :

THEOREM 3.9. Let $|k| \in \{1, \dots, n-1\}$, $A \in \mathcal{S}_0^n$, and $0 \in \text{int } A$. Then

$$\kappa_{n-1}^2 \cdot (I_k A)^\circ \subset I_k(A^\circ) \quad (3.9)$$

and equality holds if and only if A is a centered ball.

Proof. Let $X := I_k A$ and $Y := I_k(A^\circ)$. Then (3.9) is equivalent to the condition

$$\kappa_{n-1}^2 \varrho(X^\circ, u) \leq \varrho(Y, u) \quad \text{for every } u \in S^{n-1},$$

which, in view of Proposition 3.3, can be rewritten as

$$\kappa_{n-1}^2 \leq \varrho_X \cdot \varrho_Y. \quad (3.10)$$

Using the definition of I_k , the Hölder inequality, and Proposition 3.3, we obtain

$$\begin{aligned} \varrho_X(u) \cdot \varrho_Y(u) &= \frac{1}{(n-1)^2} \int_{S^{n-1} \cap u^\perp} \varrho_A(v)^k d\sigma(v) \cdot \int_{S^{n-1} \cap u^\perp} \varrho_{A^\circ}(v)^k d\sigma(v) \\ &\geq \frac{1}{(n-1)^2} \left(\int_{S^{n-1} \cap u^\perp} 1 d\sigma(v) \right)^2 = \kappa_{n-1}^2. \end{aligned}$$

Thus (3.10) is satisfied.

The equality in (3.9) holds if and only if it holds in the Hölder inequality for the functions $(\varrho_A)^{k/2}$ and $(\varrho_{A^\circ})^{k/2}$; hence, by Proposition 3.3, it holds if and only if ϱ_A is constant. This completes the proof. ■

Let us observe that, in view of Proposition 3.3, if I_k is replaced by the proportional function $J_k := (\kappa_{n-1})^{-1} I_k$, then condition (3.9) can be expressed in the more elegant form

$$(J_k A)^\circ \subset J_k(A^\circ). \quad (3.9')$$

4. FINAL REMARKS

A. The Classification Problem

To every $A \in \mathcal{S}_0^n$ we assign the subset S_A of the unit sphere:

$$S_A = \{u \in S^{n-1} \mid \varrho_A(u) > 0\}. \quad (4.1)$$

THEOREM 4.1. *Let $A_1, A_2 \in \mathcal{S}_0^n$. If there exists $f \in \text{GS}(n)$ such that $f(A_1) = A_2$, then S_{A_1} is homeomorphic to S_{A_2} .*

Proof. If $S_{A_1} = S^{n-1}$, then also $S_{A_2} = S^{n-1}$ and the assertion is trivial. Let $S^{n-1} \setminus S_{A_1} \neq \emptyset$ and let $g_i: S_{A_i} \rightarrow \text{bd } A_i \setminus \{0\}$ be the central projection for $i = 1, 2$:

$$g_i(u) = \varrho_{A_i}(u) \cdot u.$$

Clearly, g_1 and g_2 are homeomorphisms. Thus, setting

$$h(u) = (g_2)^{-1} f g_1(u) \quad \text{for } u \in S_{A_1},$$

we obtain a homeomorphism $h: S_{A_1} \rightarrow S_{A_2}$, as required. ■

The following problem is open.

Problem 1. Is the existence of a homeomorphism of S_{A_1} onto S_{A_2} sufficient for A_1, A_2 to be star equivalent?

B. Topological Properties of Star Bodies

While, evidently, star bodies in the sense of [12] are homeomorphic to B^n , the elements of \mathcal{S}_0^n are retracts of B^n and thus they are absolute retracts (see [1]):

PROPOSITION 4.2. *If $A \in \mathcal{S}_0^n$, then A is an absolute retract.*

Proof. Of course, without any loss of generality we may assume that $A \subset B^n$. Let us define $r: B^n \rightarrow A$ by the formula

$$r(x) = \begin{cases} x \varrho_A(x), & \text{if } x \in B^n \setminus A, \\ x, & \text{if } x \in A. \end{cases}$$

Then r is continuous, because

$$x \varrho_A(x) = x \quad \text{for } x \in \text{bd } A \setminus \{0\},$$

and

$$x \varrho_A(x) = u \varrho_A(u) \quad \text{for } u = \frac{x}{\|x\|},$$

where $\varrho_A(u) \rightarrow 0$ if $x \rightarrow 0 \in \text{bd } A$.

Thus, r is a retraction, since $r|_A = \text{id}$. This completes the proof. ■

C. Convex Quotient Bodies

It is easy to see that, for convex bodies A, C with $0 \in C \subset A$, the quotient star body A/C generally need not be convex, but sometimes is convex; for instance, homothetic convex bodies have convex quotient bodies:

PROPOSITION 4.3. *Let $A, C \in \mathcal{K}_0^n$ and $C \subset A$. If $A = \lambda C$ for some $\lambda > 1$, then $A/C \in \mathcal{K}_0^n$.*

Proof. Let $f(x) = \lambda x$ for every $x \in \mathbb{R}^n$. Then $f \in \text{GS}(n)$; thus, by Proposition 1.2,

$$\varrho_A(\lambda x) = \varrho_{f(C)}(f(x)) = \varrho_C(x)$$

for every $x \neq 0$. Since ϱ_A is homogeneous of degree -1 , it follows that $(1/\lambda)\varrho_A(x) = \varrho_C(x)$. This, together with Proposition 2.3, implies

$$\varrho_{A/C}(x) = (\lambda - 1)\varrho_C(x).$$

Thus, $\varrho_{A/C} = \varrho_{(\lambda-1)C}$ and consequently $A/C = (\lambda-1)C$. Hence $A/C \in \mathcal{K}_0^n$. ■

Problem 2. What can be said about the pairs (A, C) in \mathcal{K}_0^n with convex A/C ?

A contribution to this problem will be the subject of a separate paper, by Tomasz Żukowski.

ACKNOWLEDGMENTS

The author is grateful to Krzysztof Rudnik and Tomasz Żukowski for their valuable remarks.

REFERENCES

1. K. Borsuk, "Theory of Retracts," Polish Scientific Publishers, Warsaw, 1967.
2. J. Dugundji, "Topology," Allyn and Bacon, Boston, 1967.
3. R. Engelking, "General Topology," Heldermann-Verlag, Berlin, 1989.
4. R. J. Gardner, "Geometric Tomography," Cambridge Univ. Press, Cambridge, UK, 1995.
5. R. J. Gardner and A. Volčič, Tomography of convex and star bodies, *Adv. Math.* **108** (1994), 367–399.
6. I. Herburt and M. Moszyńska, On metric products, *Coll. Math.* **62** (1991), 121–133.
7. E. Lutwak, Intersection bodies and dual mixed volumes, *Adv. Math.* **71** (1988), 232–261.
8. S. MacLane, "Categories for the Working Mathematician," Springer-Verlag, Berlin, 1971.
9. M. Moszyńska, On metric inverse limits, *J. London Math. Soc.* **42** (1990), 160–174.
10. M. Moszyńska, Lipschitz inverse and direct sequences, *Topology Appl.* **56** (1994), 259–275.
11. M. Moszyńska and T. Żukowski, Duality of convex bodies, *Geom. Dedicata* **58** (1995), 161–173.
12. R. Schneider, "Convex Bodies: The Brunn–Minkowski Theory," Cambridge Univ. Press, Cambridge, UK, 1993.